



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

- (12) $cB, bcA, bAB, abcC, abBC, aAC, acABC;$
 (13) $abB, abcA, cAB, bcC, acBC, aAC, bABC;$
 (14) $acB, bA, abcAB, bcC, abBC, cAC, aABC;$
 (15) $acB, abA, bcAB, bC, abcBC, aAC, cABC;$
 (16) $acB, abcA, bAB, aC, cBC, bcAC, abABC;$
 (17) $bcB, abcA, aAB, acC, abBC, bAC, cABC;$
 (18) $abcB, acA, bAB, bcC, aBC, abAC, cABC.$

But (12) has an element in common with each of the sets (7)—(18). This is also true if we start with (13),, or (18) instead of (12). Hence the fifth, sixth,, ninth sets must be chosen from (7)—(11). Inversely, no two of the sets (6)—(11) have an element in common, and hence give a solution of the problem. We have now proved that

$$N_3 = (5.9.31)8^3(7.6.4)(6.4.2)5! \div 9! = 2^{1^2}.3.5.31).$$

Theorem. *Each of the $2^{1^2}.3.5.31$ solutions of the problem may be derived by a suitable change of notation from the solution given by the nine sets (6)—(11).*

THE UNIVERSITY OF CHICAGO, June, 1904.

DEPARTMENTS.

SOLUTIONS OF PROBLEMS.

ALGEBRA.

205. Proposed by G. B.M. ZERR, A. M., Ph. D., Parsons, W. Va.

Express in the form of radicals the roots of the equation:

$$x^{15} + 15mx^{13} + 90m^2x^{11} + 275m^3x^9 + 450m^4x^7 + 378m^5x^5 + 140m^6x^3 + 15m^7x + 2r = 0.$$

I. Solution by A. H. HOLMES, Brunswick, Maine.

$$x^{15} + 15mx^{13} + 90m^2x^{11} + 275m^3x^9 + 459m^4x^7 + 378m^5x^5 + 140m^6x^3 + 15m^7x + 2r = 0.$$

$$\therefore (x^5 + 5mx^3 + 5m^2x) + 3m^5(x^5 + 5mx^3 + 5m^2x) + 2r = 0.$$

$$\text{Put } x^5 + 5mx^3 + 5m^2x = y.$$

$$\therefore y^3 + 3m^5y + 2r = 0.$$

By the well known method of solving cubics:

$$y_1 = [-r + \sqrt{(m^{15} + r^2)}]^{\frac{1}{3}} + [-r - \sqrt{(m^{15} + r^2)}]^{\frac{1}{3}} = t_1 + t_2 = R_1,$$

$$y_2 = -\frac{1}{2}(t_1 + t_2) + \frac{1}{2}\sqrt{-3(t_1 - t_2)} = R_2,$$

$$y_3 = -\frac{1}{2}(t_1 + t_2) - \frac{1}{2}\sqrt{-3(t_1 - t_2)} = R_3,$$

$$\therefore x^5 + 5mx^3 + 5m^2x = R_1 \text{ or } R_2 \text{ or } R_3.$$

Taking the first of these values,

$$r_1 = \left(\frac{R_1}{2} + \sqrt{m^5 + \frac{R_1^2}{4}} \right)^{\frac{1}{5}} + \left(\frac{R_1}{2} - \sqrt{m^5 + \frac{R_1^2}{4}} \right)^{\frac{1}{5}} = s_1 + s_2,$$

$x_2 = es_1 + e^4s_4$; $x_3 = e^2s_1 + e^3s_4$; $x_4 = e^3s_1 + e^2s_4$, and $x_5 = e^4s_1 + es_4$, when e represents the imaginary fifth root of unity.

The other ten values of x can be found by substituting R_2 and R_3 for R_1 above.

II. Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Let $x = y + z$.

$$\begin{aligned} \therefore y^{15} + z^{15} + (yz + m)[15(y^{13} + z^{13}) + (105yz + 90m)(y'' + z'') + (455y^2z^2 + 715myz + 275m^2)(y^9 + z^9) + (1365y^3z^3 + 2925my^2z^2 + 2025m^2yz + 450m^3)(y^7 + z^7) \\ + (3003y^4z^4 + 7722my^3z^3 + 7128m^2y^2z^2 + 2772m^3yz + 378m^4)(y^5 + z^5) + (5005y^5z^5 + 14300my^4z^4 + 15400m^2y^3z^3 + 7700m^3y^2z^2 + 1750m^4yz + 140m^5)(y^3 + z^3) \\ + (6435y^6z^6 + 19305my^5z^5 + 22275m^2y^4z^4 + 12375m^3y^3z^3 + 3375m^4y^2z^2 + 405m^5yz + 15m^6)(y + z)] + 2r = 0. \end{aligned}$$

$$\therefore y^{15} + z^{15} = -2r, yz = -m. \quad \text{Let } y^{15} = a, z^{15} = b.$$

$$\therefore a + b = -2r, ab = -m^{15}. \quad \therefore a \text{ and } b \text{ are the roots of } t^2 + 2rt - m^{15} = 0.$$

Let β = an imaginary fifteenth root of unity and also let $a^{\frac{1}{15}} = c, b^{\frac{1}{15}} = d$.

$$\therefore \text{The roots are } c + d, \beta c + \beta^{14}d, \beta^{14}c + \beta d, \beta^2c + \beta^{13}d, \beta^{13}c + \beta^2d, \beta^3c + \beta^{12}d, \beta^{12}c + \beta^3d, \beta^4c + \beta^{11}d, \beta^{11}c + \beta^4d, \beta^5c + \beta^{10}d, \beta^{10}c + \beta^5d, \beta^6c + \beta^9d, \beta^9c + \beta^6d, \beta^7c + \beta^8d, \beta^8c + \beta^7d.$$

Also solved by J. Scheffer.

206. Proposed by L. E. NEWCOMB, Los Gatos, Cal.

The product of a certain pair of roots of $x^4 + ax^3 + bx^2 + amx + m^2 = 0$, is equal to the product of the remaining pair.

I. Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Let x_1, x_2, x_3, x_4 be the roots. Then $x_1 + x_2 + x_3 + x_4 = -a$; $x_1x_2x_3 + x_1x_2x_4 + x_1x_3x_4 + x_2x_3x_4 = -am$; $x_1x_2x_3x_4 = m^2$.

$$\therefore m(x_1 + x_2 + x_3 + x_4) = x_1x_2(x_3 + x_4) + x_3x_4(x_1 + x_2) = -am.$$

$$\therefore x_1x_2 = x_3x_4 = m.$$

The same is true if $x_1x_3 = x_2x_4 = m, x_1x_4 = x_2x_3 = m$.

II. Solution by J. SCHEFFER, Kee Mar College, Hagerstown, Md.

From the theory of equations, we have $a\beta\gamma + a\beta\delta + a\gamma\delta + \beta\gamma\delta = -am$, or $a\beta(\gamma + \delta) + \gamma\delta(a + \beta) = -am$, but $\gamma\delta = a\beta$.

$$\therefore a\beta(a + \beta + \gamma + \delta) = -am. \quad \text{Since } a + \beta + \gamma + \delta = -a, -a\beta a = -am.$$

$$\therefore a\beta = m = \gamma\delta.$$